

Orthogonal bundles on curves and theta functions

Arnaud BEAUVILLE

Introduction

Let C be a curve of genus $g \geq 2$, G an almost simple complex Lie group, and $\mathcal{M}_G$ the moduli space of semi-stable G -bundles on C . For each component $\mathcal{M}_G^\bullet$ of $\mathcal{M}_G$, the Picard group is infinite cyclic; its positive generator $\mathcal{L}_G^\bullet$ can be described explicitly as a determinant bundle. Then a natural question, which we will address in this paper for the classical groups, is to describe the space of “generalized theta functions” $H^0(\mathcal{M}_G^\bullet, \mathcal{L}_G^\bullet)$ and the associated rational map $\varphi_G^\bullet : \mathcal{M}_G^\bullet \dashrightarrow |\mathcal{L}_G^\bullet|^*$.

The model we have in mind is the case $G = \mathrm{SL}_r$. Let J^{g-1} be the component of the Picard variety of C parameterizing line bundles of degree $g-1$; it is isomorphic to the Jacobian of C , and carries a canonical theta divisor Θ consisting of line bundles L in J^{g-1} with $H^0(C, L) \neq 0$. For a general $E \in \mathcal{M}_{\mathrm{SL}_r}$, the locus

$$\Theta_E = \{L \in J^{g-1} \mid H^0(C, E \otimes L) \neq 0\}$$

is in a natural way a divisor, which belongs to the linear system $|r\Theta|$ on J^{g-1} . We thus obtain a rational map $\vartheta : \mathcal{M}_{\mathrm{SL}_r} \dashrightarrow |r\Theta|$. The main result of [BNR] is that *there exists an isomorphism $|\mathcal{L}_{\mathrm{SL}_r}|^* \xrightarrow{\sim} |r\Theta|$ which identifies the rational maps φ_{SL_r} and ϑ* . This gives a reasonably concrete description of φ_{SL_r} , which allows to get some information on the behaviour of this map, at least for small values of r or g (see [B3] for a survey of recent results).

Let us consider now the case $G = \mathrm{SO}_r$ with $r \geq 3$. The moduli space $\mathcal{M}_{\mathrm{SO}_r}$ parametrizes vector bundles E on C of rank r and trivial determinant, with a non-degenerate quadratic form $q : \mathbf{S}^2 E \rightarrow \mathcal{O}_C$; it has two components $\mathcal{M}_{\mathrm{SO}_r}^+$ and $\mathcal{M}_{\mathrm{SO}_r}^-$. Let $\theta : \mathcal{M}_{\mathrm{SO}_r} \dashrightarrow |r\Theta|$ be the map $(E, q) \mapsto \Theta_E$. We will see that θ maps $\mathcal{M}_{\mathrm{SO}_r}^+$ and $\mathcal{M}_{\mathrm{SO}_r}^-$ into the subspaces $|r\Theta|^+$ and $|r\Theta|^-$ corresponding to even and odd theta functions respectively. Our main result is:

Theorem. — *There are canonical isomorphisms $|\mathcal{L}_{\mathrm{SO}_r}^\pm|^* \xrightarrow{\sim} |r\Theta|^\pm$ which identify $\varphi_{\mathrm{SO}_r}^\pm : \mathcal{M}_{\mathrm{SO}_r}^\pm \dashrightarrow |\mathcal{L}_{\mathrm{SO}_r}^\pm|^*$ with the map $\theta^\pm : \mathcal{M}_{\mathrm{SO}_r}^\pm \dashrightarrow |r\Theta|^\pm$ induced by θ .*

This is easily seen to be equivalent to the fact that the pull-back map $\theta^* : H^0(J^{g-1}, \mathcal{O}(r\Theta))^* \rightarrow H^0(\mathcal{M}_{\mathrm{SO}_r}, \mathcal{L}_{\mathrm{SO}_r})$ is an *isomorphism*. We will prove that it is injective by restricting to a small subvariety of $\mathcal{M}_{\mathrm{SO}_r}$ (§ 1). Then we will use the Verlinde formula (§ 2 and 3) to show that the dimensions are the same. This is somewhat artificial since it forces us for instance to treat separately the cases r

even ≥ 6 , r odd ≥ 5 , $r = 3$ and $r = 4$. It would be interesting to find a more direct proof, perhaps in the spirit of [BNR].

In the last section we consider the same question for the symplectic group. Here the theta map does not involve the Jacobian of C but the moduli space $\mathcal{N}$ of semi-stable rank 2 vector bundles on C with determinant K_C . Let $\mathcal{L}$ be the determinant bundle on $\mathcal{N}$. For (E, φ) general in $\mathcal{M}_{\mathrm{Sp}_{2r}}$, the reduced subvariety

$$\Delta_E = \{F \in \mathcal{N} \mid H^0(E \otimes F) \neq 0\}$$

is a divisor on $\mathcal{N}$, which belongs to the linear system $|\mathcal{L}^r|$; this defines a map $\mathcal{M}_{\mathrm{Sp}_{2r}} \dashrightarrow |\mathcal{L}^r|$ which should coincide, up to a canonical isomorphism, with $\varphi_{\mathrm{Sp}_{2r}}$. This is a particular case of the *strange duality* conjecture for the symplectic group, which we discuss in § 4. Unfortunately even this particular case is not known, except in a few cases that we explain below.

1. The moduli space $\mathcal{M}_{\mathrm{SO}_r}$

(1.1) Throughout the paper we fix a complex curve C of genus $g \geq 2$. For G a semi-simple complex Lie group, we denote by $\mathcal{M}_G$ the moduli space of semi-stable G -bundles on C . It is a normal projective variety, of dimension $(g-1)\dim G$. Its connected components are in one-to-one correspondence with the elements of the group $\pi_1(G)$.

(1.2) Let us consider the case $G = \mathrm{SO}_r$ ($r \geq 3$). The space $\mathcal{M}_{\mathrm{SO}_r}$ is the moduli space of (semi-stable) orthogonal bundles, that is pairs (E, q) where E is a semi-stable¹ vector bundle of rank r and trivial determinant and $q : \mathbf{S}^2 E \rightarrow \mathcal{O}_C$ a non-degenerate quadratic form. The two components $\mathcal{M}_{\mathrm{SO}_r}^+$ and $\mathcal{M}_{\mathrm{SO}_r}^-$ are distinguished by the parity of the second Stiefel-Whitney class $w_2(E, q) \in H^2(C, \mathbf{Z}/2) \cong \mathbf{Z}/2$. This class has the following property (see e.g. [Se], Thm. 2): for every theta-characteristic κ on C and orthogonal bundle $(E, q) \in \mathcal{M}_{\mathrm{SO}_r}$,

$$w_2(E, q) \equiv h^0(C, E \otimes \kappa) + rh^0(C, \kappa) \pmod{2}. \quad (1.3)$$

The involution $\iota : L \mapsto K_C \otimes L^{-1}$ of J^{g-1} preserves Θ , hence lifts to an involution of $\mathcal{O}_{J^{g-1}}(\Theta)$. We denote by $|r\Theta|^+$ and $|r\Theta|^-$ the two corresponding eigenspaces in $|r\Theta|$, and by $\theta : \mathcal{M}_{\mathrm{SO}_r} \dashrightarrow |r\Theta|$ the map $(E, q) \mapsto \Theta_E$.

Lemma 1.4.— *The rational map $\theta : \mathcal{M}_{\mathrm{SO}_r} \dashrightarrow |r\Theta|$ maps $\mathcal{M}_{\mathrm{SO}_r}^+$ in $|r\Theta|^+$ and $\mathcal{M}_{\mathrm{SO}_r}^-$ in $|r\Theta|^-$.*

¹ By [R], 4.2, an orthogonal bundle (E, q) is semi-stable if and only if the vector bundle E is semi-stable.

Proof: For any $E \in \mathcal{M}_{\text{SL}_r}$ we have $\iota^* \Theta_E = \Theta_{E^*}$, so $\theta(\mathcal{M}_{\text{SO}_r})$ is contained in the fixed locus $|r\Theta|^+ \cup |r\Theta|^-$ of ι^* . Since $\mathcal{M}_{\text{SO}_r}^\pm$ is connected, it suffices to find one element (E, q) of $\mathcal{M}_{\text{SO}_r}^+$ (resp. $\mathcal{M}_{\text{SO}_r}^-$) such that Θ_E is a divisor in $|r\Theta|^+$ (resp. $|r\Theta|^-$).

Let $\kappa \in J^{g-1}$ be an even theta-characteristic of C ; a symmetric divisor $D \in |r\Theta|$ is in $|r\Theta|^+$ (resp. $|r\Theta|^-$) if and only if $\text{mult}_\kappa(D)$ is even (resp. odd) – see [M1], § 2. Let $J[2]$ be the 2-torsion subgroup of $\text{Pic}(C)$; we take $E = \alpha_1 \oplus \dots \oplus \alpha_r$, where $\alpha_1, \dots, \alpha_r \in J[2]$ and $\sum \alpha_i = 0$. We endow E with the diagonal quadratic form q deduced from the isomorphisms $\alpha_i^2 \cong \mathcal{O}_C$. Then $\Theta_E = \Theta_{\alpha_1} + \dots + \Theta_{\alpha_r}$. By the Riemann singularity theorem the multiplicity at κ of Θ_α is $h^0(\alpha \otimes \kappa)$. Thus by (1.3)

$$\text{mult}_\kappa(\Theta_E) = \sum_i h^0(\alpha_i \otimes \kappa) = h^0(E \otimes \kappa) \equiv w_2(E, q) \pmod{2}. \blacksquare$$

(1.5) Let $\mathcal{L}_{\text{SO}_r}$ be the determinant bundle on $\mathcal{M}_{\text{SO}_r}$, that is, the pull back of $\mathcal{L}_{\text{SL}_r}$ by the map $(E, q) \mapsto E$, and let $\mathcal{L}_{\text{SO}_r}^+$ and $\mathcal{L}_{\text{SO}_r}^-$ be its restrictions to $\mathcal{M}_{\text{SO}_r}^+$ and $\mathcal{M}_{\text{SO}_r}^-$. It follows from [BLS] that for $r \neq 4$, $\mathcal{L}_{\text{SO}_r}^\pm$ generates $\text{Pic}(\mathcal{M}_{\text{SO}_r}^\pm)$.

Proposition 1.6. – *The map $\theta^* : H^0(J^{g-1}, \mathcal{O}(r\Theta))^* \longrightarrow H^0(\mathcal{M}_{\text{SO}_r}, \mathcal{L}_{\text{SO}_r})$ induced by $\theta : \mathcal{M}_{\text{SO}_r} \dashrightarrow |r\Theta|$ is an isomorphism.*

By Lemma (1.4) θ^* splits as a direct sum $(\theta^+)^* \oplus (\theta^-)^*$, where

$$(\theta^\pm)^* : (H^0(J^{g-1}, \mathcal{O}(r\Theta))^\pm)^* \longrightarrow H^0(\mathcal{M}_{\text{SO}_r}^\pm, \mathcal{L}_{\text{SO}_r}^\pm).$$

The Proposition implies that $(\theta^+)^*$ and $(\theta^-)^*$ are isomorphisms, and this is equivalent to the Theorem stated in the introduction.

Proof of the Proposition: We will show in § 3 that the Verlinde formula gives

$$\dim H^0(\mathcal{M}_{\text{SO}_r}, \mathcal{L}_{\text{SO}_r}) = \dim H^0(J^{g-1}, \mathcal{O}(r\Theta)) = r^g.$$

It is therefore sufficient to prove that θ^* is injective, or equivalently that $\theta(\mathcal{M}_{\text{SO}_r})$ spans the projective space $|r\Theta|$. We consider again the orthogonal bundles $(E, q) = \alpha_1 \oplus \dots \oplus \alpha_r$ for $\alpha_1, \dots, \alpha_r$ in $J[2]$, $\sum \alpha_i = 0$. This bundle has a theta divisor $\Theta_E = \Theta_{\alpha_1} + \dots + \Theta_{\alpha_r}$. We claim that divisors of this form span $|r\Theta|$. To prove it, let us identify J^{g-1} with the Jacobian J of C (by choosing a divisor class of degree $g-1$). For $a \in J$, the divisor Θ_a is the only element of the linear system $|\mathcal{O}_J(\Theta) \otimes \varphi(a)|$, where $\varphi : J \rightarrow \hat{J}$ is the isomorphism associated to the principal polarization of J . Therefore our assertion follows from the following easy lemma:

Lemma 1.7. — *Let A be an abelian variety, L an ample line bundle on A , $\hat{A}[2]$ the 2-torsion subgroup of $\text{Pic}(A)$. The multiplication map*

$$\sum_{\substack{\alpha_1, \dots, \alpha_r \in \hat{A}[2] \\ \alpha_1 + \dots + \alpha_r = 0}} H^0(A, L \otimes \alpha_1) \otimes \dots \otimes H^0(A, L \otimes \alpha_r) \longrightarrow H^0(A, L^r)$$

is surjective.

Proof: Let 2_A be the multiplication by 2 in A . We have canonical isomorphisms

$$H^0(A, 2_A^* L) \cong \bigoplus_{\alpha \in \hat{A}[2]} H^0(L \otimes \alpha) \quad , \quad H^0(A, 2_A^* L^r) \cong \bigoplus_{\beta \in \hat{A}[2]} H^0(L^r \otimes \beta) ;$$

through these isomorphisms the product map $m_r : H^0(A, 2_A^* L)^{\otimes r} \longrightarrow H^0(A, 2_A^* L^r)$ is the direct sum over $\beta \in \hat{A}[2]$ of the maps

$$m_r^\beta : \sum_{\substack{\alpha_1, \dots, \alpha_r \in \hat{A}[2] \\ \alpha_1 + \dots + \alpha_r = \beta}} H^0(A, L \otimes \alpha_1) \otimes \dots \otimes H^0(A, L \otimes \alpha_r) \longrightarrow H^0(A, L^r \otimes \beta) .$$

Since the line bundle $2_A^* L$ is algebraically equivalent to L^4 , the map m_r is surjective [M2], hence so is m_r^β for every β . The case $\beta = 0$ gives the lemma. ■

2. The Verlinde formula

(2.1) We keep the notation of (1.1); we denote by q the number of simple factors of the Lie algebra of G (we are mainly interested in the case $q = 1$).

To each representation $\rho : G \rightarrow \text{SL}_r$ is attached a line bundle $\mathcal{L}_\rho$ on $\mathcal{M}_G$, the pull back of the determinant bundle on $\mathcal{M}_{\text{SL}_r}$ by the morphism $\mathcal{M}_G \rightarrow \mathcal{M}_{\text{SL}_r}$ associated to ρ . The Verlinde formula expresses the dimension of $H^0(\mathcal{M}_G, \mathcal{L}_\rho^k)$, for each integer k , in the form

$$\dim H^0(\mathcal{M}_G, \mathcal{L}_\rho^k) = N_k d_\rho(G) , \quad \text{where}$$

- $d_\rho \in \mathbb{N}^q$ is the *Dynkin index* of ρ . For $q = 1$ the number d_ρ is defined and computed in [D], § 2. In the general case the universal cover of G is a product $G_1 \times \dots \times G_q$ of almost simple factors, and we put $d_\rho = (d_{\rho_1}, \dots, d_{\rho_q})$, where ρ_i is the pull back of ρ to G_i .

We will need only to know that the Dynkin index is 2 for the standard representation of SO_r ($r \geq 5$), 4 for that of SO_3 , and $(2, 2)$ for that of SO_4 .

- $N_\ell(G)$ is an integer depending on G , the genus g of C , and $\ell \in \mathbb{N}^q$. We will now explain how this number is computed. Our basic reference is [AMW].

(2.2) *The simply connected case*

Let us first consider the case where G is simply connected and almost simple (that is, $q = 1$). Let T be a maximal torus of G , and $R = R(G, T)$ the corresponding root system (we view the roots of G as characters of T). We denote by T_ℓ the (finite) subgroup of elements $t \in T$ such that $\alpha(t) = 1$ for each *long* root α , and by T_ℓ^{reg} the subset of regular elements $t \in T_\ell$ (that is, such that $\alpha(t) \neq 1$ for each root α). It is stable under the action of the Weyl group W . For $t \in T$, we put $\Delta(t) = \prod_{\alpha \in R} (\alpha(t) - 1)$. Then the Verlinde formula is

$$N_\ell(G) = \sum_{t \in T_\ell^{\text{reg}}/W} \left(\frac{|T_\ell|}{\Delta(t)} \right)^{g-1}.$$

(2.3) This number can be explicitly computed in the following way. Let $\mathfrak{t}$ be the Lie algebra of T . The character group $P(R)$ of T embeds naturally into $\mathfrak{t}^*$. We endow $\mathfrak{t}^*$ with the W -invariant bilinear form $(\mid)$ such that $(\alpha \mid \alpha) = 2$ for each long root α , and we use this product to identify $\mathfrak{t}^*$ with $\mathfrak{t}$. Let θ be the highest root of R ; we denote by P_ℓ the set of dominant weights $\lambda \in P(R)$ such that $(\lambda \mid \theta) \leq \ell$. Let $\rho \in P(R)$ be the half-sum of the positive roots. The number $h := (\rho \mid \theta) + 1$ is the *dual Coxeter number* of R . We have $|T_\ell| = (\ell + h)^s f \nu$, where s is the rank of R , f the order of the center of G , and ν a number depending on R ; it is equal to 1 for R of type D_s and to 2 for B_s ([B2], 9.9).

For $\lambda \in P_\ell$ we put $t_\lambda = \exp 2\pi i \frac{\lambda + \rho}{\ell + h}$. The map $\lambda \mapsto t_\lambda$ is a bijection of P_ℓ onto T_ℓ^{reg}/W ([B2], 9.3.c)). For $\lambda \in P_\ell$, we have $\alpha(t_\lambda) = \exp 2\pi i \frac{(\alpha \mid \lambda + \rho)}{\ell + h}$, and therefore

$$\Delta(t_\lambda) = \prod_{\alpha \in R_+} 4 \sin^2 \pi \frac{(\alpha \mid \lambda + \rho)}{\ell + h}.$$

(2.4) *The non-simply connected case*

We now give the formula for a general almost simple group, following [AMW].

Let Z be the center of G . An element t of T belongs to Z if and only if $\alpha(t) = 1$ for all $\alpha \in R$, or equivalently $w(t) = t$ for all $w \in W$. It follows that Z acts on the set T_ℓ^{reg} by multiplication; this action commutes with that of W and thus defines an action of Z on T_ℓ^{reg}/W . Through the bijection $P_\ell \rightarrow T_\ell^{\text{reg}}/W$ the action of Z on P_ℓ is the one deduced from its action on the extended Dynkin diagram (see [O-W], § 3 or [Bo], 2.3 and 4.3).

Now let Γ be a subgroup of Z , and let $G' = G/\Gamma$. We denote by P'_ℓ the sublattice of weights $\lambda \in P_\ell$ such that $\lambda|_\Gamma = 1$. The action of Γ on P_ℓ preserves

P'_ℓ ; we denote by $\Gamma \cdot \lambda$ the orbit of a weight λ in P'_ℓ . The Verlinde formula for G' is:

$$N_\ell(G') = |\Gamma| \sum_{\lambda \in P'_\ell} |\Gamma \cdot \lambda|^{-2g} \left(\frac{|T_\ell|}{\Delta(t_\lambda)} \right)^{g-1}.$$

Each term in the sum is invariant under Γ , so we may as well sum over P'_ℓ/Γ provided we multiply each term by $|\Gamma \cdot \lambda|$:

$$N_\ell(G') = |\Gamma| \sum_{\lambda \in P'_\ell/\Gamma} |\Gamma \cdot \lambda|^{1-2g} \left(\frac{|T_\ell|}{\Delta(t_\lambda)} \right)^{g-1}. \quad (2.5)$$

(2.6) *The general case*

The above formula actually applies to any semi-simple group $G' = G/\Gamma$, where G is a product of simply connected groups $G_1, \dots, G_q$ [AMW].

We choose a maximal torus $T^{(i)}$ in G_i for each i and put $T = T^{(1)} \times \dots \times T^{(q)}$. Let $\ell := (\ell_1, \dots, \ell_q)$ be a q -uple of nonnegative integers. We put $T_\ell = T_{\ell_1}^{(1)} \times \dots \times T_{\ell_q}^{(q)}$; the subset T_ℓ^{reg} of regular elements in T_ℓ is the product of the subsets $(T_{\ell_i}^{(i)})^{\text{reg}}$. For each i , let $P_{\ell_i}^{(i)}$ be the set of dominant weights of $T^{(i)}$ associated to G_i and ℓ_i as in (2.3), and let $P_\ell = P_{\ell_1}^{(1)} \times \dots \times P_{\ell_q}^{(q)}$. For $\lambda = (\lambda_1, \dots, \lambda_q) \in P_\ell$, we put $t_\lambda = (t_{\lambda_1}, \dots, t_{\lambda_q}) \in T_\ell$; this defines a bijection of P_ℓ onto T_ℓ^{reg}/W . The elements of P_ℓ are characters of T , and we denote by P'_ℓ the subset of characters which are trivial on Γ . The group Γ is contained in the center $Z_1 \times \dots \times Z_q$ of G , which acts naturally on P_ℓ and P'_ℓ . Then

$$N_\ell(G') = |\Gamma| \sum_{\lambda \in P'_\ell/\Gamma} |\Gamma \cdot \lambda|^{1-2g} \left(\frac{|T_\ell|}{\Delta(t_\lambda)} \right)^{g-1} \quad (2.7)$$

with
$$\frac{|T_\ell|}{\Delta(t_\lambda)} = \prod_{i=1}^q \frac{|T_{\ell_i}|}{\Delta_i(t_{\lambda_i})}, \quad \Delta_i(t) = \prod_{\alpha \in R(G_i, T^{(i)})} (\alpha(t) - 1) \text{ for } t \in T^{(i)}.$$

3. The Verlinde formula for SO_r

We now apply the previous formulas to the case $G' = \text{SO}_r$. We will rest very much on the computations of [O-W]. We will borrow their notation as well as that of [Bo].

(3.1) *The case $G' = \text{SO}_{2s}$, $s \geq 3$*

The root system R is of type D_s . Let $(\varepsilon_1, \dots, \varepsilon_s)$ be the standard basis of $\mathbf{R}^s$. The weight lattice $P(R)$ is spanned by the fundamental weights

$$\varpi_j = \varepsilon_1 + \dots + \varepsilon_j \quad (1 \leq j \leq s-2),$$

$$\varpi_{s-1} = \frac{1}{2}(\varepsilon_1 + \dots + \varepsilon_{s-1} - \varepsilon_s), \quad \varpi_s = \frac{1}{2}(\varepsilon_1 + \dots + \varepsilon_{s-1} + \varepsilon_s).$$

For $\lambda \in P(R)$, we write $\lambda + \rho = \sum_i t_i \varpi_i = \sum_i u_i \varepsilon_i$ with

$$\begin{aligned} u_1 &= t_1 + \dots + t_{s-2} + \frac{1}{2}(t_{s-1} + t_s) \quad , \quad \dots \quad , \quad u_{s-2} = t_{s-2} + \frac{1}{2}(t_{s-1} + t_s) \\ u_{s-1} &= \frac{1}{2}(t_{s-1} + t_s) \quad , \quad u_s = \frac{1}{2}(-t_{s-1} + t_s) \\ (u_i &\in \frac{1}{2}\mathbf{Z} \quad , \quad u_i - u_{i+1} \in \mathbf{Z}) \quad . \end{aligned}$$

Put $k = \ell + 2s - 2$. The condition $\lambda \in P_\ell$ becomes: $u_1 > \dots > u_s$, $u_1 + u_2 < k$ and $u_{s-1} + u_s > 0$; the condition $\lambda \in P'_\ell$ imposes moreover $t_{s-1} \equiv t_s \pmod{2}$, that is, $u_i \in \mathbf{Z}$ for each i . Thus we find a bijection between P'_ℓ and the subsets $U = \{u_1, \dots, u_s\}$ of $\mathbf{Z}$ satisfying the above conditions.

The group Z is canonically isomorphic to $P(R)/Q(R)$ (note that $R = R^\vee$ in this case); its nonzero elements are the classes of ϖ_1, ϖ_{s-1} and ϖ_s . The nonzero element γ which vanishes in SO_{2s} is represented by the only weight in this list which comes from SO_{2s} , namely ϖ_1 . It corresponds to the automorphism of the extended Dynkin diagram which exchanges α_0 with α_1 and α_{s-1} with α_s (see [Bo], Table D_l); it acts on P_ℓ by $\gamma(u_1, \dots, u_s) = (k - u_1, u_2, \dots, u_{s-1}, -u_s)$. Thus the subsets U as above with $u_s \geq 0$, and moreover $u_1 \leq \frac{k}{2}$ if $u_s = 0$, form a system of representatives of P'_ℓ/Γ . The corresponding orbit has one element if $u_1 = \frac{k}{2}$ and $u_s = 0$, and 2 otherwise.

For a subset U corresponding to the weight λ we have [O-W]

$$\Delta(t_\lambda) = \Pi_k(U) = \prod_{1 \leq i < j \leq s} 4 \sin^2 \frac{\pi}{k}(u_i - u_j) 4 \sin^2 \frac{\pi}{k}(u_i + u_j) \quad .$$

Now we restrict ourselves to the case $\ell = 2$, so that $k = r = 2s$. Put $V = \{s, s-1, \dots, 0\}$. The subsets U to consider are those of the form $U_j := V - \{j\}$ for $0 \leq j \leq s$. We have

- $\Pi_r(U_j) = 4r^{s-1}$ for $1 \leq j \leq s-1$ by Corollary 1.7 (ii) in [O-W];
- $\Pi_r(U_0) = \Pi_r(U_s) = r^{s-1}$ by Corollary 1.7 (iii) in [O-W].

We have $|T_2| = 4r^s$ (2.3). Multiplying the terms U_0 and U_s by 2^{1-2g} and summing, we find:

$$N_2(SO_{2s}) = 2[(s-1).r^{g-1} + 2^{1-2g}[2.(4r)^{g-1}] = r^g \quad .$$

(3.2) *The case $G' = SO_{2s+1}$, $s \geq 2$*

Then R is of type B_s . Denoting again by $(\varepsilon_1, \dots, \varepsilon_s)$ the standard basis of $\mathbf{R}^s$, the weight lattice $P(R)$ is spanned by the fundamental weights

$$\varpi_1 = \varepsilon_1 \quad , \quad \varpi_2 = \varepsilon_1 + \varepsilon_2 \quad , \quad \dots \quad , \quad \varpi_{s-1} = \varepsilon_1 + \dots + \varepsilon_{s-1} \quad , \quad \varpi_s = \frac{1}{2}(\varepsilon_1 + \dots + \varepsilon_s) \quad .$$

For $\lambda \in P(\mathbb{R})$, we write $\lambda + \rho = \sum_i t_i \varpi_i = \sum_i u_i \varepsilon_i$ with

$$u_1 = t_1 + \dots + t_{s-1} + \frac{t_s}{2} \quad , \quad \dots \quad , \quad u_{s-1} = t_{s-1} + \frac{t_s}{2} \quad , \quad u_s = \frac{t_s}{2} \quad ,$$

with $u_i \in \frac{1}{2}\mathbb{Z}$ and $u_i - u_{i+1} \in \mathbb{Z}$ for each i . Put $k = \ell + 2s - 1$. The condition $\lambda \in P_\ell$ becomes $u_1 > \dots > u_s > 0$ and $u_1 + u_2 < k$. Since ϖ_s is the only fundamental weight which does not come from SO_{2s+1} , the condition $\lambda \in P'_\ell$ is equivalent to t_s odd, that is, $u_s \in \mathbb{Z} + \frac{1}{2}$. Thus we find a bijection between P'_ℓ and the subsets $U = \{u_1, \dots, u_s\}$ of $\mathbb{Z} + \frac{1}{2}$ satisfying

$$u_1 > \dots > u_s > 0 \quad , \quad u_1 + u_2 < k \quad .$$

The non-trivial element γ of Γ acts on P_ℓ by $\gamma(u_1, \dots, u_s) = (k - u_1, u_2, \dots, u_s)$. Thus the elements U as above with $u_1 \leq \frac{k}{2}$ form a system of representatives of P'_ℓ/Γ . The corresponding orbit has one element if $u_1 = \frac{k}{2}$ and 2 otherwise.

For a subset U corresponding to the weight λ we have [O-W]

$$\Delta(t_\lambda) = \Phi_r(U) = \prod_{1 \leq i < j \leq s} 4 \sin^2 \frac{\pi}{r} (u_i - u_j) \prod_{i=1}^s 4 \sin^2 \frac{\pi}{r} (u_i + u_j) \prod_{i=1}^s 4 \sin^2 \frac{\pi}{r} u_i \quad .$$

Now we restrict ourselves to the case $\ell = 2$, so that $k = r = 2s + 1$. Put $V = \{s + \frac{1}{2}, s - \frac{1}{2}, \dots, \frac{1}{2}\}$. The subsets U to consider are the subsets $U_j := V - \{j + \frac{1}{2}\}$ for $0 \leq j \leq s$. We have

- $\Phi_r(U_j) = 4r^{s-1}$ for $0 \leq j \leq s-1$ by Corollary 1.9 (ii) in [O-W];
- $\Phi_r(U_s) = r^{s-1}$ by Corollary 1.9 (ii) in [O-W].

We have again $|T_2| = 4r^s$ (2.3). Multiplying the term U_s by 2^{1-2g} and summing, we find:

$$N_2(\mathrm{SO}_{2s+1}) = 2[s \cdot r^{g-1} + 2^{1-2g}(4r)^{g-1}] = r^g \quad .$$

(3.3) The case $G' = \mathrm{SO}_3$

In that case $G = \mathrm{SL}_2$ has a unique fundamental weight ρ , and a unique positive root $\theta = 2\rho$. The Dynkin index of the standard representation of SO_3 is 4, so we want to compute $N_4(\mathrm{SO}_3)$. We have $|T_4| = 12$ (2.3). The set P_4 contains the weights $k\rho$ with $0 \leq k \leq 4$; the weights with k even come from SO_3 , and Γ exchanges $k\rho$ and $(4-k)\rho$. Thus a system of representatives of P'_4/Γ is $\{0, 2\rho\}$, with $|\Gamma \cdot 0| = 2$ and $|\Gamma \cdot 2\rho| = 1$. Formula (2.5) gives:

$$N_2(\mathrm{SO}_3) = 2[2^{1-2g}12^{g-1} + 3^{g-1}] = 3^g \quad .$$

(3.4) *The case $G' = \mathrm{SO}_4$*

In that case $G = \mathrm{SL}_2 \times \mathrm{SL}_2$ and the nontrivial element of Γ is $(-I, -I)$. The Dynkin index of the standard representation of SO_4 is $(2, 2)$. We have $|\mathrm{T}_2| = 8$ for SL_2 , hence $|\mathrm{T}_{(2,2)}| = 8^2$. The set $\mathrm{P}_{(2,2)}$ contains the weights $(k\rho, l\rho)$ with $0 \leq k, l \leq 2$, and $\mathrm{P}'_{(2,2)}$ is defined by the condition $k \equiv l \pmod{2}$. The element $(-I, -I)$ exchanges $(k\rho, l\rho)$ with $((2-k)\rho, (2-l)\rho)$. Thus $\mathrm{P}'_{(2,2)}/\Gamma$ consists of the classes of $(0, 0)$, $(0, 2\rho)$ and (ρ, ρ) , the latter being the only one with a nontrivial stabilizer. Formula (2.7) gives

$$N_{(2,2)}(\mathrm{SO}_4) = 2 [2 \cdot 2^{1-2g} \cdot 4^{2g-2} + 2^{2g-2}] = 4^g.$$

Therefore for each $r \geq 3$ we have obtained $\dim H^0(\mathcal{M}_{\mathrm{SO}_r}, \mathcal{L}_{\mathrm{SO}_r}) = r^g$. This achieves the proof of Proposition 1.6, and therefore of the Theorem stated in the introduction.

4. The moduli space $\mathcal{M}_{\mathrm{Sp}_{2r}}$

(4.1) Let r be an integer ≥ 1 . The space $\mathcal{M}_{\mathrm{Sp}_{2r}}$ is the moduli space of (semi-stable) symplectic bundles, that is pairs (E, φ) where E is a semi-stable² vector bundle of rank $2r$ and trivial determinant and $\varphi : \Lambda^2 E \rightarrow \mathcal{O}_C$ a non-degenerate alternate form. It is connected. To alleviate the notation we will denote it by $\mathcal{M}_r$. The determinant bundle $\mathcal{L}_r$ generates $\mathrm{Pic}(\mathcal{M}_r)$ ([K-N], [L-S]).

To describe the “strange duality” in an intrinsic way we need a variant of this space, namely the moduli space $\mathcal{M}'_r$ of semi-stable vector bundles F of rank $2r$ and determinant K_C^r , endowed with a symplectic form $\psi : \Lambda^2 F \rightarrow K_C$. If κ is a theta-characteristic on C , the map $E \mapsto E \otimes \kappa$ induces an isomorphism $\mathcal{M}_r \xrightarrow{\sim} \mathcal{M}'_r$. We denote by $\mathcal{L}'_r$ the line bundle corresponding to $\mathcal{L}_r$ under any of these isomorphisms.

Similarly, we will consider for t even the moduli space $\mathcal{M}'_{\mathrm{SO}_t}$ of semi-stable vector bundles E of rank t and determinant $K_C^{t/2}$, endowed with a quadratic form $q : \mathbf{S}^2 E \rightarrow K_C$. It has two components $\mathcal{M}'_{\mathrm{SO}_t}{}^{\pm}$ depending on the parity of $h^0(E)$; if κ is a theta-characteristic on C , the map $E \mapsto E \otimes \kappa$ induces isomorphisms $\mathcal{M}'_{\mathrm{SO}_t}{}^{\pm} \xrightarrow{\sim} \mathcal{M}'_{\mathrm{SO}_t}{}^{\pm}$ (1.3). The space $\mathcal{M}'_{\mathrm{SO}_t}{}^{+}$ carries a canonical Weil divisor, the reduced subvariety

$$\mathcal{D} = \{(E, q) \in \mathcal{M}'_{\mathrm{SO}_t}{}^{+} \mid H^0(C, E) \neq 0\};$$

$2\mathcal{D}$ is a Cartier divisor, defined by a section of the generator $\mathcal{L}'_{\mathrm{SO}_t}{}^{+}$ of $\mathrm{Pic}(\mathcal{M}'_{\mathrm{SO}_t}{}^{+})$ ([L-S], § 7).

² By the same argument as in the orthogonal case (footnote 1), a symplectic bundle (E, φ) is semi-stable if and only if E is semi-stable as a vector bundle.

(4.2) *The strange duality for symplectic bundles*

Let r, s be integers ≥ 2 , and $t = 4rs$. Consider the map

$$\pi : \mathcal{M}_r \times \mathcal{M}'_s \longrightarrow \mathcal{M}'_{\text{SO}_t}$$

which maps $((E, \varphi), (F, \psi))$ to $(E \otimes F, \varphi \otimes \psi)$. Since $\mathcal{M}_r$ is connected and contains the trivial bundle $\mathcal{O}^{2r}$ with the standard symplectic form, the image lands in $\mathcal{M}'_{\text{SO}_t}$.

For $(E, \varphi) \in \mathcal{M}_r$, the pull back of $\mathcal{L}_{\text{SO}_t}$ to $\{(E, \varphi)\} \times \mathcal{M}'_s$ is the line bundle associated to $2r$ times the standard representation, that is $\mathcal{L}_s'^{2r}$; similarly its pull back to $\mathcal{M}_r \times \{(F, \psi)\}$, for $(F, \psi) \in \mathcal{M}'_s$, is $\mathcal{L}_r^{2s}$. It follows that

$$\pi^* \mathcal{L}_{\text{SO}_t} \cong \mathcal{L}_r^{2s} \boxtimes \mathcal{L}_s'^{2r}.$$

If κ is a theta-characteristic on C with $h^0(\kappa) = 0$, we have $\pi(\mathcal{O}_C^{2r}, \kappa^{2s}) \notin \mathcal{D}$ ($\mathcal{O}_C^{2r}$ and κ^{2s} are endowed with the standard alternate forms). Thus $\Delta := \pi^* \mathcal{D}$ is a Weil divisor on $\mathcal{M}_r \times \mathcal{M}'_s$, whose double is a Cartier divisor defined by a section of $(\mathcal{L}_r^s \boxtimes \mathcal{L}_s'^r)^2$; but this moduli space is locally factorial ([S], Thm. 1.2), so that Δ is actually a Cartier divisor, defined by a section δ of $\mathcal{L}_r^s \boxtimes \mathcal{L}_s'^r$, well-defined up to a scalar. Via the Künneth isomorphism we view δ as an element of $H^0(\mathcal{M}_r, \mathcal{L}_r^s) \otimes H^0(\mathcal{M}'_s, \mathcal{L}_s'^r)$. The *strange duality* conjecture for symplectic bundles is

Conjecture 4.3.— *The section δ induces an isomorphism*

$$H^0(\mathcal{M}_r, \mathcal{L}_r^s)^* \xrightarrow{\sim} H^0(\mathcal{M}'_s, \mathcal{L}_s'^r).$$

If the conjecture holds, the rational map $\varphi_{\mathcal{L}_r^s} : \mathcal{M}_r \dashrightarrow |\mathcal{L}_r^s|$ is identified through $\delta^\#$ to the map $\mathcal{M}_r \dashrightarrow |\mathcal{L}_s'^r|$ given by $E \mapsto \Delta_E$, where Δ_E is the trace of Δ on $\{E\} \times \mathcal{M}'_s$; set-theoretically:

$$\Delta_E = \{(F, \varphi) \in \mathcal{M}'_s \mid H^0(C, E \otimes F) \neq 0\}.$$

By [O-W], we have $\dim H^0(\mathcal{M}_r, \mathcal{L}_r^s) = \dim H^0(\mathcal{M}_s, \mathcal{L}_s^r)$. Therefore the conjecture is equivalent to:

(4.4) *The linear system $|\mathcal{L}_r^{ls}|$ is spanned by the divisors Δ_E , for $E \in \mathcal{M}_r$.*

We now specialize to the case $s = 1$. The space $\mathcal{M}'_1$ is the moduli space $\mathcal{N}$ of semi-stable rank 2 vector bundles on C with determinant K_C ; its Picard group is generated by the determinant bundle $\mathcal{L}$. The conjecture becomes:

Conjecture 4.5.— *The isomorphism $\delta^\# : H^0(\mathcal{M}_r, \mathcal{L}_r)^* \xrightarrow{\sim} H^0(\mathcal{N}, \mathcal{L}^r)$ identifies the map $\varphi_{\mathcal{L}_r} : \mathcal{M}_r \dashrightarrow |\mathcal{L}_r|^*$ with the rational map $E \mapsto \Delta_E$ of $\mathcal{M}_r$ into $|\mathcal{L}|$.*

By (4.4) this is equivalent to saying that the linear system $|\mathcal{L}^r|$ on $\mathcal{N}$ is spanned by the divisors Δ_E for $E \in \mathcal{M}_r$.

(4.6) Let G be a semi-stable vector bundle of rank r and degree 0. To G is associated a divisor $\Theta_G \in |\mathcal{L}^r|$, supported on the set

$$\Theta_G = \{F \in \mathcal{N} \mid H^0(C, G \otimes F) \neq 0\}$$

provided this set is $\neq \mathcal{N}$ [D-N]. Put $E = G \oplus G^*$, with the standard symplectic form. We have $\Theta_G = \Theta_{G^*}$ by Serre duality, hence $\Delta_E = \frac{1}{2}\Theta_E = \frac{1}{2}(\Theta_G + \Theta_{G^*}) = \Theta_G$; thus conjecture 4.5 holds if the linear system $|\mathcal{L}^r|$ on $\mathcal{N}$ is spanned by the divisors Θ_G for G semi-stable of degree 0. In particular, it suffices to prove that $|\mathcal{L}^r|$ is spanned by the divisors $\Theta_{L_1} + \dots + \Theta_{L_r}$, for $L_1, \dots, L_r \in \mathcal{J}$. As a consequence of [BNR], the divisors Θ_L for L in $\mathcal{J}$ span $|\mathcal{L}|$, so *Conjecture (4.5) holds if the multiplication map $m_r : \mathbf{S}^r H^0(\mathcal{N}, \mathcal{L}) \rightarrow H^0(\mathcal{N}, \mathcal{L}^r)$ is surjective.*

Proposition 4.7.— *Conjecture 4.5 holds in the following cases:*

- (i) $r = 2$ and C has no vanishing thetanull;
- (ii) $r \geq 3g - 6$ and C is general enough;
- (iii) $g = 2$, or $g = 3$ and C is non-hyperelliptic.

Proof: In each case the multiplication map $m_r : \mathbf{S}^r H^0(\mathcal{N}, \mathcal{L}) \rightarrow H^0(\mathcal{N}, \mathcal{L}^r)$ is surjective. This follows from [B1], Prop. 2.6 c) in case (i), and from the explicit description of $\mathcal{M}_{\text{SL}_2}$ in case (iii). When C is generic, the surjectivity of m_r for r even $\geq 2g - 4$ follows from that of m_2 together with [L]. We have $H^i(\mathcal{N}, \mathcal{L}^j) = 0$ for $i \geq 1$ and $j \geq -3$ by [K-N], Thm. 2.8. By [M2] this implies that the multiplication map

$$H^0(\mathcal{N}, \mathcal{L}) \otimes H^0(\mathcal{N}, \mathcal{L}^k) \longrightarrow H^0(\mathcal{N}, \mathcal{L}^{k+1})$$

is surjective for $k \geq \dim \mathcal{N} - 3 = 3g - 6$. Together with the previous result this implies the surjectivity of m_r for $r \geq 3g - 6$, and therefore by (4.6) the Proposition. ■

Corollary 4.8.— *Suppose C has no vanishing thetanull. There is a canonical isomorphism $|\mathcal{L}_2|^* \xrightarrow{\sim} |4\Theta|^+$ which identify the maps $\varphi_{\mathcal{L}_2} : \mathcal{M}_2 \dashrightarrow |\mathcal{L}_2|^*$ with $\theta : \mathcal{M}_2 \dashrightarrow |4\Theta|^+$ such that $\theta(E, \varphi) = \Theta_E$.*

Proof: Let $i : J^{g-1} \rightarrow \mathcal{N}$ be the map $L \mapsto L \oplus \iota^* L$. The composition

$$H^0(\mathcal{M}_2, \mathcal{L}_2)^* \xrightarrow{\delta^\#} H^0(\mathcal{N}, \mathcal{L}^2) \xrightarrow{i^*} H^0(J^{g-1}, \mathcal{O}(4\Theta))^+$$

is an isomorphism by Prop. 4.7 (i) and Prop. 2.6 c) of [B1]; it maps $\varphi_{\mathcal{L}_2}(E, \varphi)$ to $i^* \Delta_E$. Using Serre duality again we find $i^* \Delta_E = \frac{1}{2}(\Theta_E + \Theta_{E^*}) = \Theta_E$, hence the Corollary. ■

(4.9) *Remarks.*— 1) The corollary does not hold if C has a vanishing theta null: the image of θ is contained in that of i^* , which is a proper subspace of $|4\Theta|^+$.

2) The analogous statement for $r \geq 3$ does not hold: the Verlinde formula implies $\dim H^0(\mathcal{N}, \mathcal{L}^r) > \dim H^0(J^{g-1}, \mathcal{O}(2r\Theta))^+$ for $g \geq 3$, or $g = 2$ and $r \geq 4$.

REFERENCES

- [AMW] A. ALEXEEV, E. MEINRENKEN, C. WOODWARD: *Formulas of Verlinde type for non-simply connected groups*. Preprint [math.SG/0005047](#).
- [B1] A. BEAUVILLE: *Fibrés de rang 2 sur les courbes, fibré déterminant et fonctions thêta II*. Bull. Soc. math. France **119** (1991), 259–291.
- [B2] A. BEAUVILLE: *Conformal blocks, Fusion rings and the Verlinde formula*. Proc. of the Hirzebruch 65 Conf. on Algebraic Geometry, Israel Math. Conf. Proc. **9** (1996), 75–96.
- [B3] A. BEAUVILLE: *Vector bundles on curves and theta functions*. Preprint [math.AG/0502179](#).
- [BLS] A. BEAUVILLE, Y. LASZLO, C. SORGER: *The Picard group of the moduli of G-bundles on a curve*. Compositio math. **112** (1998), n° 2, 183–216.
- [BNR] A. BEAUVILLE, M.S. NARASIMHAN, S. RAMANAN: *Spectral curves and the generalised theta divisor*. J. Reine Angew. Math. **398** (1989), 169–179.
- [Bo] N. BOURBAKI: *Groupes et algèbres de Lie*, Chap. VI. Hermann, Paris (1968).
- [D-N] J.-M. DREZET, M.S. NARASIMHAN: *Groupe de Picard des variétés de modules de fibrés semi-stables sur les courbes algébriques*. Invent. Math. **97** (1989), n° 1, 53–94.
- [D] E. DYNKIN: *Semisimple subalgebras of semisimple Lie algebras*. Amer. Math. Soc. Translations (II) **6** (1957), 111–244.
- [K-N] S. KUMAR, M.S. NARASIMHAN: *Picard group of the moduli spaces of G-bundles*. Math. Ann. **308** (1997), n° 1, 155–173.
- [L] Y. LASZLO: *À propos de l'espace des modules de fibrés de rang 2 sur une courbe*. Math. Ann. **299** (1994), n° 4, 597–608.
- [L-S] Y. LASZLO, C. SORGER: *The line bundles on the moduli of parabolic G-bundles over curves and their sections*. Ann. Sci. École Norm. Sup. (4) **30** (1997), n° 4, 499–525.
- [M1] D. MUMFORD: *On the equations defining abelian varieties I*. Invent. Math. **1** (1966), 287–354.
- [M2] D. MUMFORD: *Varieties defined by quadratic equations*. Questions on Algebraic Varieties (1970), 29–100; Cremonese, Rome.

- [O-W] W. OXBURY, S. WILSON: *Reciprocity laws in the Verlinde formulae for the classical groups*. Trans. Amer. Math. Soc. **348** (1996), n° 7, 2689–2710.
- [R] S. RAMANAN: *Orthogonal and spin bundles over hyperelliptic curves*. Proc. Indian Acad. Sci. Math. Sci. **90** (1981), n° 2, 151–166.
- [Se] J.-P. SERRE: *Revêtements à ramification impaire et thêta-caractéristiques*. C. R. Acad. Sci. Paris Sér. I Math. **311** (1990), n° 9, 547–552.
- [S] C. SORGER: *On moduli of G -bundles of a curve for exceptional G* . Ann. Sci. École Norm. Sup. (4) **32** (1999), n° 1, 127–133.

Arnaud BEAUVILLE
 Institut Universitaire de France
 &
 Laboratoire J.-A. Dieudonné
 UMR 6621 du CNRS
 UNIVERSITÉ DE NICE
 Parc Valrose
 F-06108 NICE Cedex 02